

Recommended Adult Immunization Schedule for ages 19 years or older

UNITED STATES

2022

How to use the adult immunization schedule

- 1** Determine recommended vaccinations by age (**Table 1**)
- 2** Assess need for additional recommended vaccinations by medical condition or other indication (**Table 2**)
- 3** Review vaccine types, frequencies, intervals, and considerations for special situations (**Notes**)
- 4** Review contraindications and precautions for vaccine types (**Appendix**)

Recommended by the Advisory Committee on Immunization Practices (www.cdc.gov/vaccines/acip) and approved by the Centers for Disease Control and Prevention (www.cdc.gov), American College of Physicians (www.acponline.org), American Academy of Family Physicians (www.aafp.org), American College of Obstetricians and Gynecologists (www.acog.org), American College of Nurse-Midwives (www.midwife.org), and American Academy of Physician Associates (www.aapa.org), and Society for Healthcare Epidemiology of America (www.shea-online.org).

Report

- Suspected cases of reportable vaccine-preventable diseases or outbreaks to the local or state health department
- Clinically significant postvaccination reactions to the Vaccine Adverse Event Reporting System at www.vaers.hhs.gov or 800-822-7967

Injury claims

All vaccines included in the adult immunization schedule except pneumococcal 23-valent polysaccharide (PPSV23) and zoster (RZV) vaccines are covered by the Vaccine Injury Compensation Program. Information on how to file a vaccine injury claim is available at www.hrsa.gov/vaccinecompensation.

Questions or comments

Contact www.cdc.gov/cdc-info or 800-CDC-INFO (800-232-4636), in English or Spanish, 8 a.m.–8 p.m. ET, Monday through Friday, excluding holidays.



Download the CDC Vaccine Schedules app for providers at www.cdc.gov/vaccines/schedules/hcp/schedule-app.html.

Helpful information

- Complete Advisory Committee on Immunization Practices (ACIP) recommendations: www.cdc.gov/vaccines/hcp/acip-recs/index.html
- *General Best Practice Guidelines for Immunization* (including contraindications and precautions): www.cdc.gov/vaccines/hcp/acip-recs/general-recs/index.html
- Vaccine information statements: www.cdc.gov/vaccines/hcp/vis/index.html
- Manual for the Surveillance of Vaccine-Preventable Diseases (including case identification and outbreak response): www.cdc.gov/vaccines/pubs/surv-manual
- Travel vaccine recommendations: www.cdc.gov/travel
- Recommended Child and Adolescent Immunization Schedule, United States, 2022: www.cdc.gov/vaccines/schedules/hcp/child-adolescent.html
- ACIP Shared Clinical Decision-Making Recommendations: www.cdc.gov/vaccines/acip/acip-scdm-faqs.html

Scan QR code for access to online schedule



CS310021-A



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention

Vaccines in the Adult Immunization Schedule*

Vaccine	Abbreviation(s)	Trade name(s)
<i>Haemophilus influenzae</i> type b vaccine	Hib	ActHIB® Hiberix® PedvaxHIB®
Hepatitis A vaccine	HepA	Havrix® Vaqta®
Hepatitis A and hepatitis B vaccine	HepA-HepB	Twinrix®
Hepatitis B vaccine	HepB	Engerix-B® Recombivax HB® Heplisav-B®
Human papillomavirus vaccine	HPV	Gardasil 9®
Influenza vaccine (inactivated)	IIV4	Many brands
Influenza vaccine (live, attenuated)	LAIV4	FluMist® Quadrivalent
Influenza vaccine (recombinant)	RIV4	Flublok® Quadrivalent
Measles, mumps, and rubella vaccine	MMR	M-M-R II®
Meningococcal serogroups A, C, W, Y vaccine	MenACWY-D MenACWY-CRM MenACWY-TT	Menactra® Menveo® MenQuadfi®
Meningococcal serogroup B vaccine	MenB-4C MenB-FHbp	Bexsero® Trumenba®
Pneumococcal 15-valent conjugate vaccine	PCV15	Vaxneuvance™
Pneumococcal 20-valent conjugate vaccine	PCV20	Prenar 20™
Pneumococcal 23-valent polysaccharide vaccine	PPSV23	Pneumovax 23®
Tetanus and diphtheria toxoids	Td	Tenivac® Tdvax™
Tetanus and diphtheria toxoids and acellular pertussis vaccine	Tdap	Adacel® Boostrix®
Varicella vaccine	VAR	Varivax®
Zoster vaccine, recombinant	RZV	Shingrix

*Administer recommended vaccines if vaccination history is incomplete or unknown. Do not restart or add doses to vaccine series if there are extended intervals between doses. The use of trade names is for identification purposes only and does not imply endorsement by the ACIP or CDC.

Table 1 Recommended Adult Immunization Schedule by Age Group, United States, 2022

Vaccine	19–26 years	27–49 years	50–64 years	≥65 years
Influenza inactivated (IIV4) or Influenza recombinant (RIV4)	1 dose annually			
Influenza live, attenuated (LAIV4)				
Tetanus, diphtheria, pertussis (Tdap or Td)	1 dose Tdap each pregnancy; 1 dose Td/Tdap for wound management (see notes)			
	1 dose Tdap, then Td or Tdap booster every 10 years			
Measles, mumps, rubella (MMR)	1 or 2 doses depending on indication (if born in 1957 or later)			
Varicella (VAR)	2 doses (if born in 1980 or later)		2 doses	
Zoster recombinant (RZV)	2 doses for immunocompromising conditions (see notes)		2 doses	
Human papillomavirus (HPV)	2 or 3 doses depending on age at initial vaccination or condition	27 through 45 years		
Pneumococcal (PCV15, PCV20, PPSV23)	1 dose PCV15 followed by PPSV23 OR 1 dose PCV20 (see notes)			1 dose PCV15 followed by PPSV23 OR 1 dose PCV20
Hepatitis A (HepA)	2 or 3 doses depending on vaccine			
Hepatitis B (HepB)	2, 3, or 4 doses depending on vaccine or condition			
Meningococcal A, C, W, Y (MenACWY)	1 or 2 doses depending on indication, see notes for booster recommendations			
Meningococcal B (MenB)	2 or 3 doses depending on vaccine and indication, see notes for booster recommendations			
	19 through 23 years			
<i>Haemophilus influenzae</i> type b (Hib)	1 or 3 doses depending on indication			

 Recommended vaccination for adults who meet age requirement, lack documentation of vaccination, or lack evidence of past infection
 Recommended vaccination for adults with an additional risk factor or another indication
 Recommended vaccination based on shared clinical decision-making
 No recommendation/Not applicable

Table 2 Recommended Adult Immunization Schedule by Medical Condition or Other Indication, United States, 2022

Vaccine	Pregnancy	Immuno-compromised (excluding HIV Infection)	HIV Infection CD4 percentage and count		Asplenia, complement deficiencies	End-stage renal disease, or on hemodialysis	Heart or lung disease; alcoholism ¹	Chronic liver disease	Diabetes	Health care personnel ²	Men who have sex with men
			<15% or <200 mm ³	≥15% and ≥200 mm ³							
IIV4 or RIV4 or LAIV4	1 dose annually										
	Contraindicated					Precaution			1 dose annually		
Tdap or Td	1 dose Tdap each pregnancy	1 dose Tdap, then Td or Tdap booster every 10 years									
MMR	Contraindicated*	Contraindicated	1 or 2 doses depending on indication								
VAR	Contraindicated*	Contraindicated		2 doses							
RZV		2 doses at age ≥19 years			2 doses at age ≥50 years						
HPV	Not Recommended*	3 doses through age 26 years			2 or 3 doses through age 26 years depending on age at initial vaccination or condition						
Pneumococcal (PCV15, PCV20, PPSV23)		1 dose PCV15 followed by PPSV23 OR 1 dose PCV20 (see notes)									
HepA				2 or 3 doses depending on vaccine							
HepB	3 doses (see notes)	2, 3, or 4 doses depending on vaccine or condition									
MenACWY		1 or 2 doses depending on indication, see notes for booster recommendations									
MenB	Precaution	2 or 3 doses depending on vaccine and indication, see notes for booster recommendations									
Hib		3 doses HSCT ³ recipients only		1 dose							

Recommended vaccination for adults who meet age requirement, lack documentation of vaccination, or lack evidence of past infection
 Recommended vaccination for adults with an additional risk factor or another indication
 Recommended vaccination based on shared clinical decision-making
 Precaution—vaccination might be indicated if benefit of protection outweighs risk of adverse reaction
 Contraindicated or not recommended—vaccine should not be administered.
 No recommendation/Not applicable

1. Precaution for LAIV4 does not apply to alcoholism. 2. See notes for influenza; hepatitis B; measles, mumps, and rubella; and varicella vaccinations. 3. Hematopoietic stem cell transplant.

Notes**Recommended Adult Immunization Schedule for ages 19 years or older, United States, 2022**

For vaccine recommendations for persons 18 years of age or younger, see the Recommended Child and Adolescent Immunization Schedule.

COVID-19 Vaccination

COVID-19 vaccines are recommended within the scope of the Emergency Use Authorization or Biologics License Application for the particular vaccine. ACIP recommendations for the use of COVID-19 vaccines can be found at www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/covid-19.html.

CDC's interim clinical considerations for use of COVID-19 vaccines can be found at www.cdc.gov/vaccines/covid-19/clinical-considerations/covid-19-vaccines-us.html.

Haemophilus influenzae type b vaccination**Special situations**

- **Anatomical or functional asplenia (including sickle cell disease):** 1 dose if previously did not receive Hib; if elective splenectomy, 1 dose, preferably at least 14 days before splenectomy
- **Hematopoietic stem cell transplant (HSCT):** 3-dose series 4 weeks apart starting 6–12 months after successful transplant, regardless of Hib vaccination history

Hepatitis A vaccination**Routine vaccination**

- **Not at risk but want protection from hepatitis A** (identification of risk factor not required): 2-dose series HepA (Havrix 6–12 months apart or Vaqta 6–18 months apart [minimum interval: 6 months]) or 3-dose series HepA-HepB (Twinrix at 0, 1, 6 months [minimum intervals: dose 1 to dose 2: 4 weeks / dose 2 to dose 3: 5 months])

Special situations

- **At risk for hepatitis A virus infection:** 2-dose series HepA or 3-dose series HepA-HepB as above
- **Chronic liver disease** (e.g., persons with hepatitis B, hepatitis C, cirrhosis, fatty liver disease, alcoholic liver disease, autoimmune hepatitis, alanine aminotransferase [ALT] or aspartate aminotransferase [AST] level greater than twice the upper limit of normal)

- **HIV infection**
- **Men who have sex with men**
- **Injection or noninjection drug use**
- **Persons experiencing homelessness**
- **Work with hepatitis A virus** in research laboratory or with nonhuman primates with hepatitis A virus infection
- **Travel in countries with high or intermediate endemic hepatitis A** (HepA-HepB [Twinrix] may be administered on an accelerated schedule of 3 doses at 0, 7, and 21–30 days, followed by a booster dose at 12 months)
- **Close, personal contact with international adoptee** (e.g., household or regular babysitting) in first 60 days after arrival from country with high or intermediate endemic hepatitis A (administer dose 1 as soon as adoption is planned, at least 2 weeks before adoptee's arrival)
- **Pregnancy** if at risk for infection or severe outcome from infection during pregnancy
- **Settings for exposure, including** health care settings targeting services to injection or noninjection drug users or group homes and nonresidential day care facilities for developmentally disabled persons (individual risk factor screening not required)

Hepatitis B vaccination**Routine vaccination**

- **Age 19 through 59 years:** complete a 2- or 3-, or 4-dose series
 - 2-dose series only applies when 2 doses of Heplisav-B* are used at least 4 weeks apart
 - 3-dose series Engerix-B or Recombivax HB at 0, 1, 6 months [minimum intervals: dose 1 to dose 2: 4 weeks / dose 2 to dose 3: 8 weeks / dose 1 to dose 3: 16 weeks]
 - 3-dose series HepA-HepB (Twinrix) at 0, 1, 6 months [minimum intervals: dose 1 to dose 2: 4 weeks / dose 2 to dose 3: 5 months]
 - 4-dose series HepA-HepB (Twinrix) accelerated schedule of 3 doses at 0, 7, and 21–30 days, followed by a booster dose at 12 months
 - 4-dose series Engerix-B at 0, 1, 2, and 6 months for persons on adult hemodialysis (note: each dosage is double that of normal adult dose, i.e., 2 mL instead of 1 mL)

***Note:** Heplisav-B not recommended in pregnancy due to lack of safety data in pregnant women

Special situations

- **Age 60 years or older* and at risk for hepatitis B virus infection:** 2-dose (Heplisav-B) or 3-dose (Engerix-B, Recombivax HB) series or 3-dose series HepA-HepB (Twinrix) as above
- **Chronic liver disease** (e.g., persons with hepatitis C, cirrhosis, fatty liver disease, alcoholic liver disease, autoimmune hepatitis, alanine aminotransferase [ALT] or aspartate aminotransferase [AST] level greater than twice upper limit of normal)
- **HIV infection**
- **Sexual exposure risk** (e.g., sex partners of hepatitis B surface antigen [HBsAg]-positive persons; sexually active persons not in mutually monogamous relationships; persons seeking evaluation or treatment for a sexually transmitted infection; men who have sex with men)
- **Current or recent injection drug use**
- **Percutaneous or mucosal risk for exposure to blood** (e.g., household contacts of HBsAg-positive persons; residents and staff of facilities for developmentally disabled persons; health care and public safety personnel with reasonably anticipated risk for exposure to blood or blood-contaminated body fluids; hemodialysis, peritoneal dialysis, home dialysis, and predialysis patients; patients with diabetes)
- **Incarcerated persons**
- **Travel in countries with high or intermediate endemic hepatitis B**

***Note:** Anyone age 60 years or older who does not meet risk-based recommendations may still receive Hepatitis B vaccination.

Human papillomavirus vaccination**Routine vaccination**

- **HPV vaccination recommended for all persons through age 26 years:** 2- or 3-dose series depending on age at initial vaccination or condition:
 - **Age 15 years or older at initial vaccination:** 3-dose series at 0, 1–2 months, 6 months (minimum intervals: dose 1 to dose 2: 4 weeks / dose 2 to dose 3: 12 weeks / dose 1 to dose 3: 5 months; repeat dose if administered too soon)
 - **Age 9–14 years at initial vaccination and received 1 dose or 2 doses less than 5 months apart:** 1 additional dose
 - **Age 9–14 years at initial vaccination and received 2 doses at least 5 months apart:** HPV vaccination series complete, no additional dose needed

Mandated NYS Infection Control Training for Healthcare Professionals

Notes Recommended Adult Immunization Schedule, United States, 2022

- **Interrupted schedules:** If vaccination schedule is interrupted, the series does not need to be restarted
- **No additional dose recommended when any HPV vaccine series has been completed using the recommended dosing intervals.**

Shared clinical decision-making

- **Some adults age 27–45 years:** Based on shared clinical decision-making, 2- or 3-dose series as above

Special situations

- **Age ranges recommended above for routine and catch-up vaccination or shared clinical decision-making also apply in special situations**
 - **Immunocompromising conditions, including HIV infection:** 3-dose series, even for those who initiate vaccination at age 9 through 14 years.
 - **Pregnancy:** Pregnancy testing is not needed before vaccination; HPV vaccination is not recommended until after pregnancy; no intervention needed if inadvertently vaccinated while pregnant

Influenza vaccination

Routine vaccination

- **Age 19 years or older:** 1 dose any influenza vaccine appropriate for age and health status annually
- For the 2021–2022 season, see www.cdc.gov/mmwr/volumes/70/rr/rr7005a1.htm
- For the 2022–23 season, see the 2022–23 ACIP influenza vaccine recommendations.

Special situations

- **Egg allergy, hives only:** any influenza vaccine appropriate for age and health status annually
- **Egg allergy—any symptom other than hives** (e.g., angioedema, respiratory distress) or required epinephrine or another emergency medical intervention: see Appendix listing contraindications and precautions
- **Severe allergic reaction (e.g., anaphylaxis) to a vaccine component or a previous dose of any influenza vaccine:** see Appendix listing contraindications and precautions
- **History of Guillain-Barré syndrome within 6 weeks after previous dose of influenza vaccine:** Generally, should not be vaccinated unless vaccination benefits outweigh risks for those at higher risk for severe complications from influenza

Measles, mumps, and rubella vaccination

Routine vaccination

- **No evidence of immunity to measles, mumps, or rubella:** 1 dose
 - **Evidence of immunity:** Born before 1957 (health care personnel, see below), documentation of receipt of MMR vaccine, laboratory evidence of immunity or disease (diagnosis of disease without laboratory confirmation is not evidence of immunity)

Special situations

- **Pregnancy with no evidence of immunity to rubella:** MMR contraindicated during pregnancy; after pregnancy (before discharge from health care facility), 1 dose
- **Nonpregnant women of childbearing age with no evidence of immunity to rubella:** 1 dose
- **HIV infection with CD4 percentages $\geq 15\%$ and CD4 count ≥ 200 cells/mm³ for at least 6 months and no evidence of immunity to measles, mumps, or rubella:** 2-dose series at least 4 weeks apart; MMR contraindicated for HIV infection with CD4 percentage $< 15\%$ or CD4 count < 200 cells/mm³
- **Severe immunocompromising conditions:** MMR contraindicated
- **Students in postsecondary educational institutions, international travelers, and household or close, personal contacts of immunocompromised persons with no evidence of immunity to measles, mumps, or rubella:** 2-dose series at least 4 weeks apart if previously did not receive any doses of MMR or 1 dose if previously received 1 dose MMR
- **Health care personnel:**
 - **Born before 1957 with no evidence of immunity to measles, mumps, or rubella:** Consider 2-dose series at least 4 weeks apart for measles or mumps or 1 dose for rubella
 - **Born in 1957 or later with no evidence of immunity to measles, mumps, or rubella:** 2-dose series at least 4 weeks apart for measles or mumps or at least 1 dose for rubella

Meningococcal vaccination

Special situations for MenACWY

- **Anatomical or functional asplenia (including sickle cell disease), HIV infection, persistent complement component deficiency, complement inhibitor (e.g., eculizumab, ravulizumab) use:** 2-dose series MenACWY-D (Menactra, Menveo, or MenQuadfi) at least 8 weeks apart and revaccinate every 5 years if risk remains
- **Travel in countries with hyperendemic or epidemic meningococcal disease, or microbiologists routinely exposed to *Neisseria meningitidis*:** 1 dose MenACWY (Menactra, Menveo, or MenQuadfi) and revaccinate every 5 years if risk remains
- **First-year college students who live in residential housing (if not previously vaccinated at age 16 years or older) or military recruits:** 1 dose MenACWY (Menactra, Menveo, or MenQuadfi)
- For MenACWY booster dose recommendations for groups listed under “Special situations” and in an outbreak setting (e.g., in community or organizational settings and among men who have sex with men) and additional meningococcal vaccination information, see www.cdc.gov/mmwr/volumes/69/rr/rr6909a1.htm

Shared clinical decision-making for MenB

- **Adolescents and young adults age 16–23 years (age 16–18 years preferred) not at increased risk for meningococcal disease:** Based on shared clinical decision-making, 2-dose series MenB-4C (Bexsero) at least 1 month apart or 2-dose series MenB-FHbp (Trumenba) at 0, 6 months (if dose 2 was administered less than 6 months after dose 1, administer dose 3 at least 4 months after dose 2); MenB-4C and MenB-FHbp are not interchangeable (use same product for all doses in series)

Special situations for MenB

- **Anatomical or functional asplenia (including sickle cell disease), persistent complement component deficiency, complement inhibitor (e.g., eculizumab, ravulizumab) use, or microbiologists routinely exposed to *Neisseria meningitidis*:**
- 2-dose primary series MenB-4C (Bexsero) at least 1 month apart or 3-dose primary series MenB-FHbp (Trumenba) at 0, 1–2, 6 months (if dose 2 was administered at least 6 months after dose 1, dose 3 not needed); MenB-4C and MenB-FHbp are not interchangeable (use same product for all doses in series); 1 dose MenB booster 1 year after primary series and revaccinate every 2–3 years if risk remains

Notes Recommended Adult Immunization Schedule, United States, 2022

- **Pregnancy:** Delay MenB until after pregnancy unless at increased risk and vaccination benefits outweigh potential risks
- For MenB **booster dose recommendations** for groups listed under “Special situations” and in an outbreak setting (e.g., in community or organizational settings and among men who have sex with men) and additional meningococcal vaccination information, see www.cdc.gov/mmwr/volumes/69/rr/rr6909a1.htm

Note: MenB vaccines may be administered simultaneously with MenACWY vaccines if indicated, but at a different anatomic site, if feasible.

Pneumococcal vaccination

Routine vaccination

- **Age 65 years or older** who have not previously received a pneumococcal conjugate vaccine or whose previous vaccination history is unknown: 1 dose PCV15 or 1 dose PCV20. If PCV15 is used, this should be followed by a dose of PPSV23 given at least 1 year after the PCV15 dose. A minimum interval of 8 weeks between PCV15 and PPSV23 can be considered for adults with an immunocompromising condition,* cochlear implant, or cerebrospinal fluid leak to minimize the risk of invasive pneumococcal disease caused by serotypes unique to PPSV23 in these vulnerable groups.
- For guidance for patients who have already received a previous dose of PCV13 and/or PPSV23, see www.cdc.gov/mmwr/volumes/71/wr/mm7104a1.htm.

Special situations

- **Age 19–64 years** with certain underlying medical conditions or other risk factors** who have not previously received a pneumococcal conjugate vaccine or whose previous vaccination history is unknown: 1 dose PCV15 or 1 dose PCV20. If PCV15 is used, this should be followed by a dose of PPSV23 given at least 1 year after the PCV15 dose. A minimum interval of 8 weeks between PCV15 and PPSV23 can be considered for adults with an immunocompromising condition,* cochlear implant, or cerebrospinal fluid leak to minimize the risk of invasive pneumococcal disease caused by serotypes unique to PPSV23 in these vulnerable groups.
- For guidance for patients who have already received a previous dose of PCV13 and/or PPSV23, see www.cdc.gov/mmwr/volumes/71/wr/mm7104a1.htm.

***Note:** Immunocompromising conditions include chronic renal failure, nephrotic syndrome, immunodeficiency, iatrogenic immunosuppression, generalized malignancy, human immunodeficiency virus, Hodgkin disease, leukemia, lymphoma, multiple myeloma, solid organ transplants, congenital or acquired asplenia, sickle cell disease, or other hemoglobinopathies.

****Note:** Underlying medical conditions or other risk factors include alcoholism, chronic heart/liver/lung disease, chronic renal failure, cigarette smoking, cochlear implant, congenital or acquired asplenia, CSF leak, diabetes mellitus, generalized malignancy, HIV, Hodgkin disease, immunodeficiency, iatrogenic immunosuppression, leukemia, lymphoma, multiple myeloma, nephrotic syndrome, solid organ transplants, or sickle cell disease or other hemoglobinopathies.

Tetanus, diphtheria, and pertussis vaccination

Routine vaccination

- **Previously did not receive Tdap at or after age 11 years:** 1 dose Tdap, then Td or Tdap every 10 years

Special situations

- **Previously did not receive primary vaccination series for tetanus, diphtheria, or pertussis:** 1 dose Tdap followed by 1 dose Td or Tdap at least 4 weeks after Tdap and another dose Td or Tdap 6–12 months after last Td or Tdap (Tdap can be substituted for any Td dose, but preferred as first dose), Td or Tdap every 10 years thereafter
- **Pregnancy:** 1 dose Tdap during each pregnancy, preferably in early part of gestational weeks 27–36
- **Wound management:** Persons with 3 or more doses of tetanus-toxoid-containing vaccine: For clean and minor wounds, administer Tdap or Td if more than 10 years since last dose of tetanus-toxoid-containing vaccine; for all other wounds, administer Tdap or Td if more than 5 years since last dose of tetanus-toxoid-containing vaccine. Tdap is preferred for persons who have not previously received Tdap or whose Tdap history is unknown. If a tetanus-toxoid-containing vaccine is indicated for a pregnant woman, use Tdap. For detailed information, see www.cdc.gov/mmwr/volumes/69/wr/mm6903a5.htm

Varicella vaccination

Routine vaccination

- **No evidence of immunity to varicella:** 2-dose series 4–8 weeks apart if previously did not receive varicella-containing vaccine (VAR or MMRV [measles-mumps-rubella-varicella vaccine] for children); if previously received 1 dose varicella-containing vaccine, 1 dose at least 4 weeks after first dose

- Evidence of immunity: U.S.-born before 1980 (except for pregnant women and health care personnel [see below]), documentation of 2 doses varicella-containing vaccine at least 4 weeks apart, diagnosis or verification of history of varicella or herpes zoster by a health care provider, laboratory evidence of immunity or disease

Special situations

- **Pregnancy with no evidence of immunity to varicella:** VAR contraindicated during pregnancy; after pregnancy (before discharge from health care facility), 1 dose if previously received 1 dose varicella-containing vaccine or dose 1 of 2-dose series (dose 2: 4–8 weeks later) if previously did not receive any varicella-containing vaccine, regardless of whether U.S.-born before 1980
- **Health care personnel with no evidence of immunity to varicella:** 1 dose if previously received 1 dose varicella-containing vaccine; 2-dose series 4–8 weeks apart if previously did not receive any varicella-containing vaccine, regardless of whether U.S.-born before 1980
- **HIV infection with CD4 percentages $\geq 15\%$ and CD4 count ≥ 200 cells/mm³ with no evidence of immunity:** Vaccination may be considered (2 doses 3 months apart); VAR contraindicated for HIV infection with CD4 percentage $< 15\%$ or CD4 count < 200 cells/mm³
- **Severe immunocompromising conditions:** VAR contraindicated

Zoster vaccination

Routine vaccination

- **Age 50 years or older:** 2-dose series RZV (Shingrix) 2–6 months apart (minimum interval: 4 weeks; repeat dose if administered too soon), regardless of previous herpes zoster or history of zoster vaccine live (ZVL, Zostavax) vaccination (administer RZV at least 2 months after ZVL)

Special situations

- **Pregnancy:** There is currently no ACIP recommendation for RZV use in pregnancy. Consider delaying RZV until after pregnancy.
- **Immunocompromising conditions (including HIV):** RZV recommended for use in persons age 19 years or older who are or will be immunodeficient or immunosuppressed because of disease or therapy. For detailed information, see www.cdc.gov/mmwr/volumes/71/wr/mm7103a2.htm.

Appendix Recommended Adult Immunization Schedule, United States, 2022

Guide to Contraindications and Precautions to Commonly Used Vaccines

Adapted from Table 4-1 in Advisory Committee on Immunization Practices (ACIP) General Best Practice Guidelines for Immunization: Contraindication and Precautions available at www.cdc.gov/vaccines/hcp/acip-recs/general-recs/contraindications.html and ACIP's Recommendations for the Prevention and Control of 2021-22 Seasonal Influenza with Vaccines available at www.cdc.gov/mmwr/volumes/70/rr/rr7005a1.htm

Interim clinical considerations for use of COVID-19 vaccines including contraindications and precautions can be found at www.cdc.gov/vaccines/covid-19/clinical-considerations/covid-19-vaccines-us.html

Vaccine	Contraindications ¹	Precautions ²
Influenza, egg-based, inactivated injectable (IIV4)	<ul style="list-style-type: none"> Severe allergic reaction (e.g., anaphylaxis) after previous dose of any influenza vaccine (i.e., any egg-based IIV, cclIV, RIV, or LAIV of any valency) Severe allergic reaction (e.g., anaphylaxis) to any vaccine component³ (excluding egg) 	<ul style="list-style-type: none"> Guillain-Barré syndrome (GBS) within 6 weeks after a previous dose of any type of influenza vaccine Persons with egg allergy with symptoms other than hives (e.g., angioedema, respiratory distress) or required epinephrine or another emergency medical intervention: Any influenza vaccine appropriate for age and health status may be administered. If using egg-based IIV4, administer in medical setting under supervision of health care provider who can recognize and manage severe allergic reactions. May consult an allergist. Moderate or severe acute illness with or without fever
Influenza, cell culture-based inactivated injectable [(cclIV4), Fluceivax* Quadrivalent]	<ul style="list-style-type: none"> Severe allergic reaction (e.g., anaphylaxis) to any cclIV of any valency, or to any component³ of cclIV4 	<ul style="list-style-type: none"> Guillain-Barré syndrome (GBS) within 6 weeks after a previous dose of any type of influenza vaccine Persons with a history of severe allergic reaction (e.g., anaphylaxis) after a previous dose of any egg-based IIV, RIV, or LAIV of any valency. If using cclIV4, administer in medical setting under supervision of health care provider who can recognize and manage severe allergic reactions. May consult an allergist. Moderate or severe acute illness with or without fever
Influenza, recombinant injectable [(RIV4), Flublok* Quadrivalent]	<ul style="list-style-type: none"> Severe allergic reaction (e.g., anaphylaxis) to any RIV of any valency, or to any component³ of RIV4 	<ul style="list-style-type: none"> Guillain-Barré syndrome (GBS) within 6 weeks after a previous dose of any type of influenza vaccine Persons with a history of severe allergic reaction (e.g., anaphylaxis) after a previous dose of any egg-based IIV, cclIV, or LAIV of any valency. If using RIV4, administer in medical setting under supervision of health care provider who can recognize and manage severe allergic reactions. May consult an allergist. Moderate or severe acute illness with or without fever
Influenza, live attenuated [LAIV4, Flumist* Quadrivalent]	<ul style="list-style-type: none"> Severe allergic reaction (e.g., anaphylaxis) after previous dose of any influenza vaccine (i.e., any egg-based IIV, cclIV, RIV, or LAIV of any valency) Severe allergic reaction (e.g., anaphylaxis) to any vaccine component³ (excluding egg) Adults age 50 years or older Anatomic or functional asplenia Immunocompromised due to any cause including, but not limited to, medications and HIV infection Close contacts or caregivers of severely immunosuppressed persons who require a protected environment Pregnancy Cochlear implant Active communication between the cerebrospinal fluid (CSF) and the oropharynx, nasopharynx, nose, ear, or any other cranial CSF leak Received influenza antiviral medications oseltamivir or zanamivir within the previous 48 hours, peramivir within the previous 5 days, or baloxavir within the previous 17 days. 	<ul style="list-style-type: none"> Guillain-Barré syndrome (GBS) within 6 weeks after a previous dose of any type of influenza vaccine Asthma in persons aged 5 years old or older Persons with egg allergy with symptoms other than hives (e.g., angioedema, respiratory distress) or required epinephrine or another emergency medical intervention: Any influenza vaccine appropriate for age and health status may be administered. If using LAIV4 (which is egg based), administer in medical setting under supervision of health care provider who can recognize and manage severe allergic reactions. May consult an allergist. Persons with underlying medical conditions (other than those listed under contraindications) that might predispose to complications after wild-type influenza virus infection [e.g., chronic pulmonary, cardiovascular (except isolated hypertension), renal, hepatic, neurologic, hematologic, or metabolic disorders (including diabetes mellitus)] Moderate or severe acute illness with or without fever

- When a contraindication is present, a vaccine should NOT be administered. Kroger A, Bahta L, Hunter P. ACIP General Best Practice Guidelines for Immunization. www.cdc.gov/vaccines/hcp/acip-recs/general-recs/contraindications.html
- When a precaution is present, vaccination should generally be deferred but might be indicated if the benefit of protection from the vaccine outweighs the risk for an adverse reaction. Kroger A, Bahta L, Hunter P. ACIP General Best Practice Guidelines for Immunization. www.cdc.gov/vaccines/hcp/acip-recs/general-recs/contraindications.html
- Vaccination providers should check FDA-approved prescribing information for the most complete and updated information, including contraindications, warnings, and precautions. Package inserts for U.S.-licensed vaccines are available at www.fda.gov/vaccines-blood-biologics/approved-products/vaccines-licensed-use-united-states.

Appendix Recommended Adult Immunization Schedule, United States, 2022

Vaccine	Contraindications ¹	Precautions ²
<i>Haemophilus influenzae</i> type b (Hib)	<ul style="list-style-type: none"> Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component³ For Hiberix, Act-Hib, and Pedvaxi-HIB only: History of severe allergic reaction to dry natural latex 	<ul style="list-style-type: none"> Moderate or severe acute illness with or without fever
Hepatitis A (HepA)	<ul style="list-style-type: none"> Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component³ including neomycin 	<ul style="list-style-type: none"> Moderate or severe acute illness with or without fever
Hepatitis B (HepB)	<ul style="list-style-type: none"> Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component³ including yeast For HepB only: Pregnancy 	<ul style="list-style-type: none"> Moderate or severe acute illness with or without fever
Hepatitis A- Hepatitis B vaccine [HepA-HepB, (Twinrix®)]	<ul style="list-style-type: none"> Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component³ including neomycin and yeast 	<ul style="list-style-type: none"> Moderate or severe acute illness with or without fever
Human papillomavirus (HPV)	<ul style="list-style-type: none"> Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component³ 	<ul style="list-style-type: none"> Moderate or severe acute illness with or without fever
Measles, mumps, rubella (MMR)	<ul style="list-style-type: none"> Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component³ Severe immunodeficiency (e.g., hematologic and solid tumors, receipt of chemotherapy, congenital immunodeficiency, long-term immunosuppressive therapy or patients with HIV infection who are severely immunocompromised) Pregnancy Family history of altered immunocompetence, unless verified clinically or by laboratory testing as immunocompetent 	<ul style="list-style-type: none"> Recent (≤11 months) receipt of antibody-containing blood product (specific interval depends on product) History of thrombocytopenia or thrombocytopenic purpura Need for tuberculin skin testing or interferon-gamma release assay (IGRA) testing Moderate or severe acute illness with or without fever
Meningococcal ACWY (MenACWY) [MenACWY-CRM (Menveo®); MenACWY-D (Menactra®); MenACWY-TT (MenQuadfi®)]	<ul style="list-style-type: none"> Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component³ For MenACWY-D and Men ACWY-CRM only: severe allergic reaction to any diphtheria toxoid- or CRM197-containing vaccine For MenACWY-TT only: severe allergic reaction to a tetanus toxoid-containing vaccine 	<ul style="list-style-type: none"> Moderate or severe acute illness with or without fever
Meningococcal B (MenB) [MenB-4C (Bexsero); MenB-FHbp (Trumenb)]	<ul style="list-style-type: none"> Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component³ 	<ul style="list-style-type: none"> Pregnancy For MenB-4C only: Latex sensitivity Moderate or severe acute illness with or without fever
Pneumococcal conjugate (PCV15)	<ul style="list-style-type: none"> Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component³ Severe allergic reaction (e.g., anaphylaxis) to any diphtheria-toxoid-containing vaccine or to its vaccine component³ 	<ul style="list-style-type: none"> Moderate or severe acute illness with or without fever
Pneumococcal conjugate (PCV20)	<ul style="list-style-type: none"> Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component³ Severe allergic reaction (e.g., anaphylaxis) to any diphtheria-toxoid-containing vaccine or to its vaccine component³ 	<ul style="list-style-type: none"> Moderate or severe acute illness with or without fever
Pneumococcal polysaccharide (PPSV23)	<ul style="list-style-type: none"> Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component³ 	<ul style="list-style-type: none"> Moderate or severe acute illness with or without fever
Tetanus, diphtheria, and acellular pertussis (Tdap) Tetanus, diphtheria (Td)	<ul style="list-style-type: none"> Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component³ For Tdap only: Encephalopathy (e.g., coma, decreased level of consciousness, prolonged seizures), not attributable to another identifiable cause, within 7 days of administration of previous dose of DTP, DTaP, or Tdap 	<ul style="list-style-type: none"> Guillain-Barré syndrome (GBS) within 6 weeks after a previous dose of tetanus-toxoid-containing vaccine History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria-toxoid-containing or tetanus-toxoid-containing vaccine; defer vaccination until at least 10 years have elapsed since the last tetanus-toxoid-containing vaccine Moderate or severe acute illness with or without fever For Tdap only: Progressive or unstable neurological disorder, uncontrolled seizures, or progressive encephalopathy until a treatment regimen has been established and the condition has stabilized
Varicella (VAR)	<ul style="list-style-type: none"> Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component³ Severe immunodeficiency (e.g., hematologic and solid tumors, receipt of chemotherapy, congenital immunodeficiency, long-term immunosuppressive therapy or patients with HIV infection who are severely immunocompromised) Pregnancy Family history of altered immunocompetence, unless verified clinically or by laboratory testing as immunocompetent 	<ul style="list-style-type: none"> Recent (≤11 months) receipt of antibody-containing blood product (specific interval depends on product) Receipt of specific antiviral drugs (acyclovir, famciclovir, or valacyclovir) 24 hours before vaccination (avoid use of these antiviral drugs for 14 days after vaccination) Use of aspirin or aspirin-containing products Moderate or severe acute illness with or without fever
Zoster recombinant vaccine (RZV)	<ul style="list-style-type: none"> Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component³ 	<ul style="list-style-type: none"> Moderate or severe acute illness with or without fever Current herpes zoster infection

- When a contraindication is present, a vaccine should NOT be administered. Kroger A, Bahta L, Hunter P. ACIP General Best Practice Guidelines for Immunization. www.cdc.gov/vaccines/hcp/acip-recs/general-recs/contraindications.html
- When a precaution is present, vaccination should generally be deferred but might be indicated if the benefit of protection from the vaccine outweighs the risk for an adverse reaction. Kroger A, Bahta L, Hunter P. ACIP General Best Practice Guidelines for Immunization. www.cdc.gov/vaccines/hcp/acip-recs/general-recs/contraindications.html
- Vaccination providers should check FDA-approved prescribing information for the most complete and updated information, including contraindications, warnings, and precautions. Package inserts for U.S.-licensed vaccines are available at www.fda.gov/vaccines-blood-biologics/approved-products/vaccines-licensed-use-united-states.

New York State Department of Health

Adult Immunization Schedule

Vaccine/Age	19-26 Years	27-59 Years	60-64 Years	65 Years or Older	
Influenza (Flu)	You need one dose every year.				
Tetanus, Diphtheria, Pertussis (Td, Tdap)	<p>There are two vaccines that are commonly referred to as the “tetanus shot”: Td (tetanus and diphtheria) and Tdap (tetanus, diphtheria, and pertussis). Adults need both vaccines at different times during their lives.</p> <p>If you are pregnant, you need one dose of Tdap between 27 and 36 weeks of pregnancy, preferably closer to 27 weeks. You will need a repeat dose every time you are pregnant to protect yourself and your baby.</p> <p>If you’ve never had, or don’t remember having, a tetanus shot before, you will need one Tdap shot followed by two Td shots. You will need a Td shot every 10 years.</p> <p>Be sure to consult your health care provider if you have a deep or dirty wound. You may need a dose at that time.</p>				
Zoster (Shingles)			You need one dose of this vaccine if you are 60 or older.		
Pneumococcal (PCV13, PPSV23)	<p>There are two vaccines that protect against pneumococcal disease: PCV13 and PPSV23. Adults may need both vaccines at different times in their lives.</p> <p>You need one or more doses of PPSV23 if you smoke cigarettes or have chronic health conditions.*</p> <p>People with certain medical conditions, including weak immune systems, may also need a dose of PCV13.*</p>			You need both PCV13 and PPSV23 at 65 or older.	
Varicella (Chickenpox)	<p>If you’ve never had chickenpox, or you were never vaccinated, you need to get two doses.</p> <p>If you’ve been vaccinated but you received only one dose, you need to get the second dose as soon as possible.</p>				
Human Papillomavirus (HPV)	<p>You need three doses if you are a woman who is 26 or younger, or a man who is 21 or younger and did not already complete the series. You might only need two doses if you started as a teenager. Some men aged 22-26 should also get the HPV vaccine.*</p>				
Measles, Mumps, Rubella (MMR)	<p>You may need one or two doses if you were born in 1957 or later and you were never vaccinated.*</p> <p>You may need one or two doses if you were born before 1957 and you work in the health care field.*</p>				

*Consult your health care provider to determine your risk for infection and your need for this vaccine.

Mandated NYS Infection Control Training for Healthcare Professionals

New York State Department of Health

Adult Immunization Schedule

Vaccine/Age	19-26 Years	27-59 Years	60-64 Years	65 Years or Older
Meningococcal	<p>There are two vaccines that protect against meningococcal disease: MCV4 (meningococcal conjugate) and MenB (serogroup B meningococcal). You need one or more doses of MCV4 vaccine if you:</p> <ul style="list-style-type: none"> • are a first-year college student who is 21 or younger living in communal housing, • are a military recruit, • live or work in certain other high-risk settings, or • have certain medical conditions.* <p>Ask your health care provider about MenB vaccine if you do not have a spleen (or it does not work well), have sickle cell disease, have complement component deficiency or are 23 or younger.*</p>			
Hepatitis A	You need two doses if you have never been fully vaccinated and you are either at risk for hepatitis A* or wish to be protected.			
Hepatitis B	You need three doses if you have never been fully vaccinated and you are either at risk for hepatitis B* or wish to be protected.			
<i>Haemophilus influenzae type b (Hib)</i>	You need one or more doses of this vaccine if you do not have a spleen (or it doesn't work well), have sickle cell disease, or received a bone marrow transplant.*			
Do you travel outside the United States?	<p>If yes, you may need some of the above, and additional, vaccines. Please consult with your health care provider or a travel health specialist. For more information about vaccine recommendations for your travel destination, you can call the Centers for Disease Control and Prevention (CDC) at 1-800-232-4636, or visit their website at www.cdc.gov/travel.</p>			

*Consult your health care provider to determine your risk for infection and your need for this vaccine.



health.ny.gov
 facebook.com/NYSDOH
 twitter.com/HealthNYGov

New York State Recommended Childhood and Adolescent Immunization Schedule

A check ✓ means that this is the earliest and best time for your child to be immunized. If your child misses the “best time” for vaccination, he or she should still be immunized as quickly as possible. Ask your doctor about getting your child caught up.

Vaccine against:	Birth	2 months	4 months	6 months	12 months	15 months	18-23 months	4-6 years	11-12 years	16 years	
Hepatitis B	✓	✓ 1-2 mo.		✓ 6-18 mo.							
Rotavirus		✓	✓	✓ ¹							
Diphtheria, Tetanus, Pertussis (DTaP)		✓	✓	✓		✓ 15-18 mo.		✓			
Tetanus, Diphtheria, Pertussis (Tdap) ²									✓ ²		
<i>Haemophilus influenzae</i> type b (Hib)		✓	✓	✓ ¹	✓ 12-15 mo.						
Pneumococcal Disease (PCV) ³		✓	✓	✓	✓ 12-15 mo.		Ask your doctor if your child 2 years old or older should get vaccinated with PPSV23. ³				
Polio (IPV)		✓	✓	✓ 6-18 mo.				✓			
Influenza				Recommended yearly for all children aged 6 months and older. Ask your doctor if your child should receive one or two doses.							
Measles, Mumps, Rubella (MMR) ⁴				See footnote 4	✓ 12-15 mo.			✓			
Varicella (Chickenpox)					✓ 12-15 mo.			✓			
Hepatitis A					✓		✓				
Human Papillomavirus (HPV) ⁵									✓ ⁵		
Meningococcal Disease ⁶		Ask your doctor if your child 2 months old or older should get vaccinated against meningococcal disease.							✓		✓

¹ For some types of Hib and Rotavirus vaccine, the 6-month dose is not needed.

² Tdap: Children 7-10 years old who are not fully immunized against pertussis should receive a single dose of Tdap.

³ PCV = Pneumococcal Conjugate Vaccine; PPSV23 = Pneumococcal Polysaccharide Vaccine

⁴ MMR: Children 6-11 months old who are traveling outside the U.S. should receive one dose of MMR before departure.

⁵ The HPV vaccine includes two shots given 6 months apart. It is recommended for both boys and girls. Teens who start the series after age 15, and some children with special medical conditions, may need three doses.

⁶ There are two vaccines that protect against meningococcal disease. Some children with special medical conditions may need both MCV4 and MenB.